


PSJ2 Exh 48



Pain Management

May 5, 2009

Dr. Ira Kornbluth MD
President Maryland Pain Society

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Intro to Pain Management

- Types of pain
- Assessment of pain
- Pharmacological treatment options
- Non-Pharmacological treatment options

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Epidemiology of Pain

- An estimated 76.5 million Americans (26%) aged 20 years and over report that they have had a problem with pain of any sort that persisted for more than 24 hours in duration (not including acute pain)
- 20% of American adults (42 million people) report that pain or physical discomfort disrupts their sleep a few nights a week or more
- Chronic pain disables more people than cancer or heart disease and costs the American people more than both combined
- The annual cost of chronic pain in the US, including healthcare expenses, lost income, and lost productivity is estimated to be \$100 billion

Source: APF, Decision Resources- Chronic Pain, March 2008; National Center for Health Statistics Report: Health, US, 2006, Special Feature on Pain



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



Epidemiology of Pain

Disease Population in the US	% Diagnosed	% Drug Treated
Osteoarthritis pain	53	50
Rheumatoid arthritis pain	98	95
Cancer pain		
<i>Severe</i>	85	75
<i>Bone</i>	75	75
<i>Neuropathic</i>	70	60
<i>Breakthrough</i>	80	75
Chronic low back pain	65	80
Fibromyalgia	22	85
HIV-related neuropathic pain	38	53

Source: APF: Decision Resources- Chronic Pain, March 2008; National Center for Health Statistics Report: Health, US, 2006, Special Feature on Pain

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

What is Pain?

- **Medical definition**

“Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage.”

» International Association for the Study of Pain, 1979
- **Patient-Derived Definition**


“Pain is whatever the experiencing person says it is, existing whenever he/she says it does”.

» Margo McCaffrey

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS



Unresolved chronic pain has a negative impact on the sufferers' quality of life, affecting ability to concentrate, do their job, exercise, socialize, get a good night's sleep, perform household chores, and have sex. It affects their participation in activities, results in missed work, and is associated with fatigue, frustration, anxiety, irritability, and depression. Pain interferes with their ability to sleep, productivity, and ability to work. Some feel they cannot function as normal people and feel so bad they want to die.

Reference

American Pain Foundation. Pain facts. Found at:
www.painfoundation.org/page.asp?file=Library/PainSurveys.htm. Accessed on
 October 19, 2006.

Disability


- 86 million Americans suffer from chronic pain
- Low Back Pain
 - Leading cause of disability in adults <45
 - 3rd leading cause of disability in adults >45
 - Nearly 8 million people in US partially disabled due to LBP

Sources:
http://www.theacpa.org/nerve/about_acpa.asp
<http://emedicine.medscape.com/article/310353-overview>
http://www.karger.com/gazette/65/walsh/art_4_p.htm

THE STRENGTH WITHIN


CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS



Return to Work

- Less than 50% if off work for greater than 6 months
- 25% will return if > 1 year
- Nearly 0% if > 2 years




Waddell G. The clinical course of low back pain. In: *The back pain revolution*. Edinburgh: Churchill Livingstone, 1998:103-117.

THE STRENGTH WITHIN


CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS



Pain Assessment


- Character
- Onset
- Location
- Duration
- Intensity
- Alleviating factors
- Exacerbating factors



THE STRENGTH WITHIN


CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS



HNP/bulge on MRI

- Jensen found 64% of adults without back pain had a bulge, protrusion, or extrusion at one level
- 38% had abnormality at more than one level





THE
STRENGTH
WITHIN

Jensen MC et al. Magnetic resonance imaging of the lumbar spine in people without back pain. N Engl J Med 1994;331:69-73

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS







Portenoy RK, Kanner RM. Definition and Assessment of Pain. In: Portenoy RK, Kanner RM, eds. *Pain Management: Theory and Practice*. Philadelphia, Pa: FA Davis Company; 1996:4.

Galer BS, Dworkin RH. *A Clinical Guide to Neuropathic Pain*. Minneapolis, Minn: The McGraw-Hill Companies Inc; 2000:8-9.

Nociceptive Pain

Somatic vs. Visceral

- **Somatic**
 - Deep or cutaneous tissue injury; localized
 - Described as dull, achy, gnawing
 - Responsive to opioids & NSAIDs
 - Examples
 - Tumor, bone mets, fractures, muscle pulls
- **Visceral**
 - Organ system origin; poorly localized
 - Described as diffuse and achy
 - Responsive to opioids
 - Examples
 - Tumor infiltrate, bowel obstruction, pancreatic CA



THE STRENGTH WITHIN

For Training Purposes Only
Not Intended for Promotion

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS


Neuropathic Pain

- Results from damage to or changes of peripheral or central nervous system
- Often described as burning, sharp, electrical, lightning symptoms
- Peripheral Neuropathies
 - Diabetic, Alcohol.
- Entrapment Neuropathies
 - CTS, Neuromas
- CRPS
 - RSD
- Central Pain
 - Post CVA Pain, MS




CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed





5th Vital Sign

1. Temperature
2. Pulse
3. Respiration rate
4. Blood pressure
5. PAIN



The illustration depicts a medical examination. A doctor in a white coat is shown from the side, examining a patient who is seated and facing away. The doctor is holding a stethoscope to the patient's back. Above the doctor, a thermometer is shown with a radiating line, indicating temperature. To the right, a blood pressure cuff is shown with a clock face, indicating pulse or blood pressure. Below the doctor, a small inset shows a person holding their arm, indicating pain. To the right of the inset, a large pair of lungs is shown, indicating respiration rate.

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

What Are the Goals of Clinical Assessment?

- Achieve diagnosis of pain and underlying disorder
- Identify pain mechanism
- Evaluate functional status (ADLs*)
- Identify comorbid conditions
- Evaluate psychosocial factors
- Set goals
- Develop a targeted treatment plan
- Determine when to refer

*ADLs= activities of daily living



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



Pain Assessment

- Pain History

- Onset
- Location
- Duration
- Radiation
- Exacerbating factors
- Alleviating factors
- Describe pain quality
- Rate pain severity
- Affect on
 - Sleep
 - Relationships
 - ADOL


- Pain treatment history
- History of substance abuse
- Physical Exam
- Diagnostic Studies



ADOL = activities of daily living

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed






Pain Measurement Tools

- Pain Diary
- Visual Analog Scale (VAS)
- Verbal Numerical Scale
- Wisconsin Brief Pain Inventory (BPI)
- McGill Pain Questionnaire
- WOMAC Osteoarthritis Index
- Neuropathic Pain Scale (NPS)



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed






Katz N. *Clin J Pain*. 2000;16:S41-S48;
Belgrade MJ. *Postgrad Med*. 1999;106:127-148;
Kim KM et al. *J Korean Med Sci*. 2001;16:193-197;
Galer BS et al. *A Clinical Guide to Neuropathic Pain*. 2000;11,90;
Turk DC. *Clin J Pain*. 2000;16:279-280.

24


Nonpharmacologic Options

- Biofeedback
- Relaxation therapy
- Physical and occupational therapy
- Cognitive/behavioral strategies
 - meditation; guided imagery
- Acupuncture
- Transcutaneous electrical nerve stimulation (TENS)




For Training Purposes Only
Not Intended for Promotion

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



Ferrell B, Herr K, Epplin J, et al. The management of persistent pain in older persons. *Programs and Abstracts of the American Geriatric Society 2002 Annual Scientific Meeting*, May 8–12, 2002: Washington, DC

Kuman D, Marshall HJ. Diabetic peripheral neuropathy: amelioration of pain with transcutaneous electrostimulation. *Diabetes Care*. 1997;20:1702-1705.



1. Mackin GA. *J Hand Ther.* 1997;10:96-109.
2. Katz N. *Clin J Pain.* 2000;16:S41-S48.
3. Leland JY. *Geriatrics.* 1999;54:23-37.
4. Belgrade MJ. *Postgrad Med.* 1999;106:127-140.
5. Galer BS et al. *A Clinical Guide to Neuropathic Pain.* 2000:97.
6. Gonzales GR. *Neurology.* 1995;45(suppl 9):S11-S16.

Medication Management

- Reversible, titratable
- Adjuvant meds treat concurrent symptoms
- Use of multiple drugs is typical in chronic pain mgmt
- Lack of invasiveness



THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDOT
PHARMACEUTICALS

Limitations of Pharmacotherapy

- Abuse, diversion
- Compliance
- Side effects
- If pain is localized to one region, is there a targeted treatment (e.g. injection, surgery)?




THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Opioid Treatment of Acute and Chronic Pain

- Use of analgesics should include
 - Proper patient selection
 1. A proper initial assessment of the patient
 2. The use of opioids as part of a comprehensive treatment plan e.g., consider non-opioid treatments as well
 3. Patient reassessment- analgesia? Adverse effects? Activities of daily living? Aberrant behaviors?
 4. Opioid rotation- rotation from one opioid to another can result in significant improvement in analgesia



THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Overview of Current Opioid Environment

- Dichotomy currently exists:
 - Increased awareness of pain & need to treat
 - Opioids recognized as effective analgesics
 - Abuse/addiction/diversion issues raise concerns
 - Result of this dichotomy = barriers to treatment
- Need for a “balanced” approach
 - Appropriate access to pain medications
 - Proactive efforts to limit abuse, addiction, diversion



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



Opioid Basics

- No ceiling dose
- No End Organ Toxicity
- Opioid tolerance will develop!!!!
- Opioid dependence may develop.
- Opioid addiction unlikely to develop.
 - If pain exists then addiction = 1: 10,000

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Controlled Substance Act

- Scheduling due to medical use & potential for abuse
- Five schedules: C-I, C-II, C-III, C-IV, C-V
- C-I: No accepted medical use (i.e., LSD)
- C-II: Accepted med. use; high abuse potential
 - i.e., Morphine, hydromorphone, oxymorphone
- C-III: “Less” abuse vs. C-1 or C-2
 - i.e., Hydrocodone w/APAP

APAP=Acetaminophen

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Definition Related to Opioid Use

- **Physical Dependence**
 - Defined solely by the occurrence of an abstinence syndrome on abrupt dose reduction or discontinuation.
- **Tolerance**
 - Diminution of one or more drug effects caused by exposure to drug
- **Addiction**
 - Aberrant use of substance characterized by loss of control, compulsive use, and continued use despite harm to self



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



	OPANA ER	OxyContin	Avinza	Kadian
Active Compound	Oxymorphone	Oxycodone	Morphine	Morphine
Dosing	Pivotal trials demonstrate proven every 12 hour dosing that is maintained over 12 week period	Every 12 hours	Once-a-day	Once or twice-a-day
Half-life	IR = up to 9 hours ER = up to 11.3 hours	IR = 3.2 hours CR = 4.5 hours	Not listed	Not listed
Metabolism	Metabolized in the liver. No CYP450 drug/drug interactions at clinically relevant doses. No dose adjustment necessary for CYP3A4 or CYP 2C9-mediated interactions	CYP450 2D6	Conjugated in the liver	Conjugated in the liver
Relative Potency	10 mg	20 mg	30 mg	30 mg
Delivery System	TIMERx-N matrix	AcroContin - biphasic	Encapsulated beads	Encapsulated pellets



THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



ENDO
PHARMACEUTICALS

34

OPANA[®] ER Indication

think 
OPANA[®] ER
(oxymorphone HCl) 
Extended-release tablets
5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg

- OPANA ER is indicated for the relief of moderate to severe pain in patients requiring continuous, around-the-clock opioid treatment for an extended period of time
- OPANA ER is not intended for use as an as needed analgesic
- OPANA ER is not indicated for pain in the immediate post-operative period (12–24 hours following surgery) for patients not previously taking opioids because of the risk of oversedation and respiratory depression requiring reversal with opioid antagonists
- OPANA ER is not indicated for pain in the post-operative period if the pain is mild or not expected to persist for an extended period of time


 CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed 

35


OPANA® ER

Proven Durable Efficacy

- OPANA ER has demonstrated proven every 12-hour dosing
 - Patented TIMERx®-N extended-release delivers steady release of oxymorphone over the entire 12-hour dosing period
- Durable analgesic effect and significant pain relief demonstrated in opioid-naïve and opioid-experienced patients
 - Minimal increase in pain scores over a 12-week period
 - Patients successfully titrated to an effective dose
 - Maintained effective analgesic dose over the 12-week period
- Less rescue medicine use than the allowable daily maximum
 - In clinical trials, patients were allowed up to 10 mg/day of OPANA®, an immediate-release formulation
 - Average use was consistently less than the allowed 10 mg/day (7.3 mg–9.8 mg for opioid-naïve pts and 7.5 mg–8.5 mg for opioid-experienced pts)



CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed



36

Dosing Advantages with OPANA® ER


- No known CYP450 PK drug-drug interactions at clinically relevant doses
 - Multiple concomitant medications may be a significant issue for some patients
 - Many common drugs are processed via the CYP450 pathway, therefore, no dose adjustments are required for these drugs
- Clinically studied in over 2,000 patients including studies in low back pain, osteoarthritis, and cancer pain
 - Opioid-naïve patients and opioid-experienced patients
- Multiple strengths, multiple formulations of oxymorphone
 - Ease of initiation of therapy
 - Ease of conversion within the OPANA® brand family
 - Flexibility when converting from other opioids

THE STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

ENDO
PHARMACEUTICALS

Chronic Pain Outcome Measures



- Reduction in pain
- Reduced consumption of analgesics
- Enhanced ADLs
- Return to work
- Other functional ability outcomes

ADL = Activities of Daily Living

STRENGTH WITHIN

CONFIDENTIAL- for internal training purposes only.
Not to be used in promotion with customers. Not to be distributed

END
PHARMACEUTICALS

- The outcome measures I have³ used to organize the results reported in the literature are reduction in pain, discontinued or greatly reduced consumption of oral analgesics, enhanced performance of activities of daily living, return to work and other functional ability outcomes.

